

Overview: Physically-Based Simulation of Single Event Effects

Mike Alles
Scooter Ball
Dan Fleetwood
Christina Howe
Aaron Kobayashi
Lloyd Massengill

Marcus Mendenhall
Ron Schrimpf
Andrew Sternberg
Robert Reed
Bob Weller


MURI Annual Review 2004
Vanderbilt University
16/August/2004

Supported in part by DTRA, NASA, the U.S. Air Force, and the DOD MFEL program.





<http://www.f22-raptor.com/>




[http://www.icknowledge.com/trends/p4\(2\).jpg](http://www.icknowledge.com/trends/p4(2).jpg)


<http://www.pbs.org/wgbh/buildingbig/>




Physically-Based SEE Simulation



Physically-Based SEE Simulation



Physically-Based SEE Simulation




Overview

- What is our strategy?
- What is GEANT4?
- Why use it?
- How is the strategy implemented?
- What are the results so far?
 - Delta electron single events.
 - Displacement single events.
- What is next?



GEANT4? A Simulation Toolkit...

S. Agostinelli et al. "Geant4 - a simulation toolkit," Nucl. Instr. Meth. A506 (2003) 250.



100 MeV protons \rightarrow 5 μ m Si

VANDERBILT UNIVERSITY


Why use GEANT4 for radiation effects in electronics?

For an excited system, the average response, is not, in general, the response to the average excitation.

Translation: For very small devices, all events are *single* events.






Implementation? How it works...




Result: MOS transistor response to energetic δ -ray events

Supply = 3.3 V $R_L = 200 \text{ k}\Omega$ Biased off




Drain Current Pulses

100 MeV protons




Load line = $R_L = 200 \text{ k}\Omega$




The Shape of Large-Pulse Events


Most energetic event: 48.7 keV



35.8 keV
1.0 μA




37.6 keV
1.2 μA




41.3 keV
1.0 μA



A large event : $\approx 10\% V_{DS}$



Particle Trajectories
Incident proton: blue
Electrons: red



Energy in the transistor: 44.7 keV




Result: A displacement damage module for Geant4

- Objective: Add screened Coulomb scattering to Geant4
- Why?
- Incorporate TRIM/SRIM functionality
- Compute nonionizing energy deposition to estimate displacement damage
- Low priority for Geant4 collaboration



Validation: Rutherford Scattering

2 MeV α particles on 100 nm of Si




Conclusion: Large angle scattering is correct




Validation: Ion Implantation

Boron in Si



Arsenic in Si




Conclusion: Energy deposition processes are correct




Validation: Multiple Scattering

270 keV projectiles on $100 \mu\text{g}/\text{cm}^2$ of C

α particles




Protons




Conclusion: Small-angle collisions are correct




Displacement Damage in GaAs



Viewing Displacement Damage




A Displacement Single Event



How does energy translate to electrical activity?



Physically-Based SEE Simulation



Summary

- The basic framework for ensemble simulation of single events is in place and tested.
- The methodology has been demonstrated by application to low-probability, high-energy, discrete δ -ray events.
- A screened Coulomb scattering module has been written and tested for displacement damage computations.
- Our first event tool for rectangular samples is begins testing this week...



What's next?

- Incorporate realistic device geometry
- Simulate proton, heavy-ion nuclear reactions
- Develop SEE cross section estimates
- Compute systematic nonionizing energy values
- Analyze devices - small, new technologies
- Couple energy deposition/defect theory
- Develop derivative models, software engineering
- *Validation, validation, validation!*



^{56}Fe on Si: 56 GeV

